

Journal of Inequalities in Pure and Applied Mathematics


NEIGHBORHOODS OF CERTAIN CLASSES OF ANALYTIC FUNCTIONS OF COMPLEX ORDER

G. MURUGUSUNDARAMOORTHY AND H. M. SRIVASTAVA

Department of Mathematics
Vellore Institute of Technology (Deemed University)
Vellore 632014, Tamil Nadu
India
EMail: gmsmoorthy@rediffmail.com

Department of Mathematics and Statistics
University of Victoria
Victoria, British Columbia V8W 3P4
Canada
EMail: harimsri@math.uvic.ca

©2000 [Victoria University](#)
ISSN (electronic): 1443-5756
024-04



volume 5, issue 2, article 24,
2004.

*Received 30 January, 2004;
accepted 02 February, 2004.*

Communicated by: Th.M. Rassias

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract


By making use of the familiar concept of neighborhoods of analytic functions, the authors prove several inclusion relations associated with the (n, δ) -neighborhoods of certain subclasses of analytic functions of complex order, which are introduced here by means of the Ruscheweyh derivatives. Special cases of some of these inclusion relations are shown to yield known results.

2000 Mathematics Subject Classification: Primary 30C45.

Key words: Analytic functions, Ruscheweyh derivatives, Starlike functions, Convex functions, δ -neighborhood, Inclusion relations.

Contents

1	Introduction and Definitions	3
2	Inclusion Relations Involving $N_{n,\delta}(e)$	7
3	Neighborhoods for the Classes $\mathcal{S}_n^{(\alpha)}(\gamma, \lambda, \beta)$ and $\mathcal{R}_n^{(\alpha)}(\gamma, \lambda, \beta; \mu)$	13
	References	



Neighborhoods of Certain Classes of Analytic Functions of Complex Order

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back

Close

Quit

Page 2 of 18

1. Introduction and Definitions

Let $\mathcal{A}(n)$ denote the class of functions f of the form:

$$(1.1) \quad f(z) = z - \sum_{k=n+1}^{\infty} a_k z^k \quad (a_k \geq 0; k \in \mathbb{N} \setminus \{1\}; n \in \mathbb{N} := \{1, 2, 3, \dots\}),$$

which are *analytic* in the *open unit disk*

$$\mathbb{U} := \{z : z \in \mathbb{C} \text{ and } |z| < 1\}.$$

Following the works of Goodman [9] and Ruscheweyh [14], we define the (n, δ) -neighborhood of a function $f \in \mathcal{A}(n)$ by (see also [2], [3], [4], and [16])


$$(1.2) \quad N_{n,\delta}(f) := \left\{ g \in \mathcal{A}(n) : g(z) = z - \sum_{k=n+1}^{\infty} b_k z^k \right. \\ \left. \text{and } \sum_{k=n+1}^{\infty} k |a_k - b_k| \leq \delta \right\}.$$

In particular, for the *identity* function

$$(1.3) \quad e(z) = z,$$

we immediately have

$$(1.4) \quad N_{n,\delta}(e) := \left\{ g \in \mathcal{A}(n) : g(z) = z - \sum_{k=n+1}^{\infty} b_k z^k \text{ and } \sum_{k=n+1}^{\infty} k |b_k| \leq \delta \right\}.$$



**Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order**

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back

Close

Quit

Page 3 of 18

The above concept of (n, δ) -neighborhoods was extended and applied recently to families of *analytically* multivalent functions by Altıntaş *et al.* [6] and to families of *meromorphically* multivalent functions by Liu and Srivastava ([10] and [11]). The main object of the present paper is to investigate the (n, δ) -neighborhoods of several subclasses of the class $\mathcal{A}(n)$ of *normalized* analytic functions in \mathbb{U} with negative *and* missing coefficients, which are introduced below by making use of the Ruscheweyh derivatives.

First of all, we say that a function $f \in \mathcal{A}(n)$ is *starlike of complex order* γ ($\gamma \in \mathbb{C} \setminus \{0\}$), that is, $f \in \mathcal{S}_n^*(\gamma)$, if it also satisfies the following inequality:

$$(1.5) \quad \Re \left(1 + \frac{1}{\gamma} \left[\frac{zf'(z)}{f(z)} - 1 \right] \right) > 0 \quad (z \in \mathbb{U}; \gamma \in \mathbb{C} \setminus \{0\}).$$


Furthermore, a function $f \in \mathcal{A}(n)$ is said to be *convex of complex order* γ ($\gamma \in \mathbb{C} \setminus \{0\}$), that is, $f \in \mathcal{C}_n(\gamma)$, if it also satisfies the following inequality:

$$(1.6) \quad \Re \left(1 + \frac{1}{\gamma} \frac{zf''(z)}{f'(z)} \right) > 0 \quad (z \in \mathbb{U}; \gamma \in \mathbb{C} \setminus \{0\}).$$

The classes $\mathcal{S}_n^*(\gamma)$ and $\mathcal{C}_n(\gamma)$ stem essentially from the classes of starlike and convex functions of complex order, which were considered earlier by Nasr and Aouf [12] and Wiatrowski [18], respectively (see also [5] and [7]).

Next, for the functions f_j ($j = 1, 2$) given by

$$(1.7) \quad f_j(z) = z + \sum_{k=2}^{\infty} a_{k,j} z^k \quad (j = 1, 2),$$



**Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order**

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back

Close

Quit

Page 4 of 18

let $f_1 * f_2$ denote the Hadamard product (or convolution) of f_1 and f_2 , defined by

$$(1.8) \quad (f_1 * f_2)(z) := z + \sum_{k=2}^{\infty} a_{k,1} a_{k,2} z^k =: (f_2 * f_1)(z).$$

Thus the Ruscheweyh derivative operator $D^\lambda : \mathcal{A} \rightarrow \mathcal{A}$ is defined for $\mathcal{A} := \mathcal{A}(1)$ by

$$(1.9) \quad D^\lambda f(z) := \frac{z}{(1-z)^{\lambda+1}} * f(z) \quad (\lambda > -1; f \in \mathcal{A})$$

or, equivalently, by


$$(1.10) \quad D^\lambda f(z) := z - \sum_{k=2}^{\infty} \binom{\lambda + k - 1}{k - 1} a_k z^k \quad (\lambda > -1; f \in \mathcal{A})$$

for a function $f \in \mathcal{A}$ of the form (1.1). Here, *and in what follows*, we make use of the following standard notation:

$$(1.11) \quad \binom{\kappa}{k} := \frac{\kappa(\kappa - 1) \cdots (\kappa - k + 1)}{k!} \quad (\kappa \in \mathbb{C}; k \in \mathbb{N}_0)$$

for a binomial coefficient. In particular, we have

$$(1.12) \quad D^n f(z) = \frac{z(z^{n-1} f(z))^{(n)}}{n!} \quad (n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}).$$



**Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order**

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back

Close

Quit

Page 5 of 18

Finally, in terms of the Ruscheweyh derivative $D^\lambda (\lambda > -1)$ defined by (1.9) or (1.10) above, let $\mathcal{S}_n(\gamma, \lambda, \beta)$ denote the subclass of $\mathcal{A}(n)$ consisting of functions f which satisfy the following inequality:

$$(1.13) \quad \left| \frac{1}{\gamma} \left(\frac{z (D^\lambda f(z))'}{D^\lambda f(z)} - 1 \right) \right| < \beta$$

$$(z \in \mathbb{U}; \gamma \in \mathbb{C} \setminus \{0\}; \lambda > -1; 0 < \beta \leq 1).$$

Also let $\mathcal{R}_n(\gamma, \lambda, \beta; \mu)$ denote the subclass of $\mathcal{A}(n)$ consisting of functions f which satisfy the following inequality:


$$(1.14) \quad \left| \frac{1}{\gamma} \left((1 - \mu) \frac{D^\lambda f(z)}{z} + \mu (D^\lambda f(z))' - 1 \right) \right| < \beta$$

$$(z \in \mathbb{U}; \gamma \in \mathbb{C} \setminus \{0\}; \lambda > -1; 0 < \beta \leq 1; 0 \leq \mu \leq 1).$$

Various further subclasses of the classes $\mathcal{S}_n(\gamma, \lambda, \beta)$ and $\mathcal{R}_n(\gamma, \lambda, \beta; \mu)$ with $\gamma = 1$ were studied in many earlier works (cf., e.g., [8] and [17]; see also the references cited in these earlier works). Clearly, in the case of (for example) the class $\mathcal{S}_n(\gamma, \lambda, \beta)$, we have

$$(1.15) \quad \mathcal{S}_n(\gamma, 0, 1) \subset \mathcal{S}_n^*(\gamma) \quad \text{and} \quad \mathcal{S}_n(\gamma, 1, 1) \subset \mathcal{C}_n(\gamma)$$

$$(n \in \mathbb{N}; \gamma \in \mathbb{C} \setminus \{0\}).$$



Neighborhoods of Certain Classes of Analytic Functions of Complex Order

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back

Close

Quit

Page 6 of 18

2. Inclusion Relations Involving $N_{n,\delta}(e)$

In our investigation of the inclusion relations involving $N_{n,\delta}(e)$, we shall require Lemma 1 and Lemma 2 below.

Lemma 1. *Let the function $f \in \mathcal{A}(n)$ be defined by (1.1). Then f is in the class $\mathcal{S}_n(\gamma, \lambda, \beta)$ if and only if*

$$(2.1) \quad \sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} (\beta|\gamma| + k - 1) a_k \leq \beta|\gamma|.$$


Proof. We first suppose that $f \in \mathcal{S}_n(\gamma, \lambda, \beta)$. Then, by appealing to the condition (1.13), we readily obtain

$$(2.2) \quad \Re \left(\frac{z (D^\lambda f(z))'}{D^\lambda f(z)} - 1 \right) > -\beta|\gamma| \quad (z \in \mathbb{U})$$

or, equivalently,

$$(2.3) \quad \Re \left(\frac{-\sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} (k-1) a_k z^k}{z - \sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} a_k z^k} \right) > -\beta|\gamma| \quad (z \in \mathbb{U}),$$

where we have made use of (1.10) and the definition (1.1). We now choose values of z on the real axis and let $z \rightarrow 1-$ through real values. Then the



Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back

Close

Quit

Page 7 of 18

inequality (2.3) immediately yields the desired condition (2.1). Conversely, by applying the hypothesis (2.1) and letting $|z| = 1$, we find that

$$\begin{aligned}
 \left| \frac{z (D^\lambda f(z))'}{D^\lambda f(z)} - 1 \right| &= \left| \frac{\sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} (k-1) a_k z^k}{z - \sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} a_k z^k} \right| \\
 &\leq \frac{\beta |\gamma| \left(1 - \sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} a_k \right)}{1 - \sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} a_k} \\
 (2.4) \qquad \qquad \qquad &\leq \beta |\gamma|.
 \end{aligned}$$

Hence, by the maximum modulus theorem, we have


$$f \in \mathcal{S}_n(\gamma, \lambda, \beta),$$

which evidently completes the proof of Lemma 1. □

Similarly, we can prove the following result.

Lemma 2. *Let the function $f \in \mathcal{A}(n)$ be defined by (1.1). Then f is in the class $\mathcal{R}(\gamma, \lambda, \beta; \mu)$ if and only if*

$$(2.5) \qquad \sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} [\mu(k-1) + 1] a_k \leq \beta |\gamma|.$$



**Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order**

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back

Close

Quit

Page 8 of 18

Remark 1. A special case of Lemma 1 when

$$n = 1, \quad \gamma = 1, \quad \text{and} \quad \beta = 1 - \alpha \quad (0 \leq \alpha < 1)$$

was given earlier by Ahuja [1]. Furthermore, if in Lemma 1 with

$$n = 1, \quad \gamma = 1, \quad \text{and} \quad \beta = 1 - \alpha \quad (0 \leq \alpha < 1),$$

we set $\lambda = 0$ and $\lambda = 1$, we shall obtain the familiar results of Silverman [15].

Our first inclusion relation involving $N_{n,\delta}(e)$ is given by Theorem 1 below.

Theorem 1. If

$$(2.6) \quad \delta := \frac{(n+1)\beta|\gamma|}{(\beta|\gamma|+n)\binom{\lambda+n}{n}} \quad (|\gamma| < 1),$$

then


$$(2.7) \quad \mathcal{S}_n(\gamma, \lambda, \beta) \subset N_{n,\delta}(e).$$

Proof. For a function $f \in \mathcal{S}_n(\gamma, \lambda, \beta)$ of the form (1.1), Lemma 1 immediately yields

$$(\beta|\gamma|+n)\binom{\lambda+n}{n} \sum_{k=n+1}^{\infty} a_k \leq \beta|\gamma|,$$

so that

$$(2.8) \quad \sum_{k=n+1}^{\infty} a_k \leq \frac{\beta|\gamma|}{(\beta|\gamma|+n)\binom{\lambda+n}{n}}.$$



**Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order**

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back

Close

Quit

Page 9 of 18

On the other hand, we also find from (2.1) and (2.8) that

$$\begin{aligned} \binom{\lambda+n}{n} \sum_{k=n+1}^{\infty} k a_k &\leq \beta|\gamma| + (1-\beta|\gamma|) \binom{\lambda+n}{n} \sum_{k=n+1}^{\infty} a_k \\ &\leq \beta|\gamma| + (1-\beta|\gamma|) \binom{\lambda+n}{n} \frac{\beta|\gamma|}{(\beta|\gamma|+n) \binom{\lambda+n}{n}} \\ &\leq \frac{(n+1)\beta|\gamma|}{\beta|\gamma|+n} \quad (|\gamma| < 1), \end{aligned}$$

that is,

$$(2.9) \quad \sum_{k=n+1}^{\infty} k a_k \leq \frac{(n+1)\beta|\gamma|}{(\beta|\gamma|+n) \binom{\lambda+n}{n}} := \delta,$$

which, in view of the definition (1.4), proves Theorem 1. □


By similarly applying Lemma 2 instead of Lemma 1, we now prove Theorem 2 below.

Theorem 2. *If*

$$(2.10) \quad \delta := \frac{(n+1)\beta|\gamma|}{(\mu n+1) \binom{\lambda+n}{n}},$$

then

$$(2.11) \quad \mathcal{R}_n(\gamma, \lambda, \beta; \mu) \subset N_{n,\delta}(e).$$



**Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order**

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back

Close

Quit

Page 10 of 18

Proof. Suppose that a function $f \in \mathcal{R}(\gamma, \lambda, \beta; \mu)$ is of the form (1.1). Then we find from the assertion (2.5) of Lemma 2 that

$$\binom{\lambda + n}{n} (\mu n + 1) \sum_{k=n+1}^{\infty} a_k \leq \beta |\gamma|,$$

which yields the following coefficient inequality:

$$(2.12) \quad \sum_{k=n+1}^{\infty} a_k \leq \frac{\beta |\gamma|}{(\mu n + 1) \binom{\lambda + n}{n}}.$$

Making use of (2.5) in conjunction with (2.12), we also have

$$(2.13) \quad \begin{aligned} \mu \binom{\lambda + n}{n} \sum_{k=n+1}^{\infty} k a_k &\leq \beta |\gamma| + (\mu - 1) \binom{\lambda + n}{n} \sum_{k=n+1}^{\infty} a_k \\ &\leq \beta |\gamma| + (\mu - 1) \binom{\lambda + n}{n} \frac{\beta |\gamma|}{(\mu n + 1) \binom{\lambda + n}{n}}, \end{aligned}$$

that is,

$$\sum_{k=n+1}^{\infty} k a_k \leq \frac{(n + 1) \beta |\gamma|}{(\mu n + 1) \binom{\lambda + n}{n}} =: \delta,$$

which, in light of the definition (1.4), completes the proof of Theorem 2. \square



**Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order**

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents




Go Back

Close

Quit

Page 11 of 18

Remark 2. By suitably specializing the various parameters involved in Theorem 1 and Theorem 2, we can derive the corresponding inclusion relations for many relatively more familiar function classes (see also Equation (1.15) and Remark 1 above).



**Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order**

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back

Close

Quit

Page 12 of 18

3. Neighborhoods for the Classes $\mathcal{S}_n^{(\alpha)}(\gamma, \lambda, \beta)$ and $\mathcal{R}_n^{(\alpha)}(\gamma, \lambda, \beta; \mu)$

In this section we determine the neighborhood for each of the classes

$$\mathcal{S}_n^{(\alpha)}(\gamma, \lambda, \beta) \quad \text{and} \quad \mathcal{R}_n^{(\alpha)}(\gamma, \lambda, \beta; \mu),$$

which we define as follows. A function $f \in \mathcal{A}(n)$ is said to be in the class $\mathcal{S}_n^{(\alpha)}(\gamma, \lambda, \beta)$ if there exists a function $g \in \mathcal{S}_n(\gamma, \lambda, \beta)$ such that

$$(3.1) \quad \left| \frac{f(z)}{g(z)} - 1 \right| < 1 - \alpha \quad (z \in \mathbb{U}; 0 \leq \alpha < 1).$$


Analogously, a function $f \in \mathcal{A}(n)$ is said to be in the class $\mathcal{R}_n^{(\alpha)}(\gamma, \lambda, \beta; \mu)$ if there exists a function $g \in \mathcal{R}_n(\gamma, \lambda, \beta; \mu)$ such that the inequality (3.1) holds true.

Theorem 3. *If $g \in \mathcal{S}_n(\gamma, \lambda, \beta)$ and*

$$(3.2) \quad \alpha = 1 - \frac{(\beta |\gamma| + n) \delta \binom{\lambda + n}{n}}{(n + 1) \left[(\beta |\gamma| + n) \binom{\lambda + n}{n} - \beta |\gamma| \right]},$$

then

$$(3.3) \quad N_{n, \delta}(g) \subset \mathcal{S}_n^{(\alpha)}(\gamma, \lambda, \beta).$$



Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back

Close

Quit

Page 13 of 18

Proof. Suppose that $f \in N_{n,\delta}(g)$. We then find from the definition (1.2) that

$$(3.4) \quad \sum_{k=n+1}^{\infty} k |a_k - b_k| \leq \delta,$$

which readily implies the coefficient inequality:


$$(3.5) \quad \sum_{k=n+1}^{\infty} |a_k - b_k| \leq \frac{\delta}{n+1} \quad (n \in \mathbb{N}).$$

Next, since $g \in \mathcal{S}_n(\gamma, \lambda, \beta)$, we have [cf. Equation (2.8)]

$$(3.6) \quad \sum_{k=n+1}^{\infty} b_k \leq \frac{\beta|\gamma|}{(\beta|\gamma| + n) \binom{\lambda+n}{n}},$$

so that

$$(3.7) \quad \begin{aligned} \left| \frac{f(z)}{g(z)} - 1 \right| &< \frac{\sum_{k=n+1}^{\infty} |a_k - b_k|}{1 - \sum_{k=n+1}^{\infty} b_k} \\ &\leq \frac{\delta}{n+1} \cdot \frac{(\beta|\gamma| + n) \binom{\lambda+n}{n}}{(\beta|\gamma| + n) \binom{\lambda+n}{n} - \beta|\gamma|} \\ &= 1 - \alpha, \end{aligned}$$



**Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order**

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back

Close

Quit

Page 14 of 18

provided that α is given precisely by (3.2). Thus, by definition, $f \in \mathcal{S}_n^{(\alpha)}(\gamma, \lambda, \beta)$ for α given by (3.2). This evidently completes our proof of Theorem 3. \square

Our proof of Theorem 4 below is much akin to that of Theorem 3.


Theorem 4. *If $g \in \mathcal{R}_n(\gamma, \lambda, \beta; \mu)$ and*

$$(3.8) \quad \alpha = 1 - \frac{(\mu n + 1) \delta \binom{\lambda + n}{n}}{(n + 1) \left[(\mu n + 1) \binom{\lambda + n}{n} - \beta |\gamma| \right]},$$

then

$$(3.9) \quad N_{n,\delta}(g) \subset \mathcal{R}_n^{(\alpha)}(\gamma, \lambda, \beta; \mu).$$

Remark 3. *Just as we indicated already in Section 1 and Remark 2, Theorem 3 and Theorem 4 can readily be specialized to deduce the corresponding neighborhood results for many simpler function classes.*



**Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order**

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back


Close

Quit

Page 15 of 18

References

- [1] O.P. AHUJA, Hadamard products of analytic functions defined by Ruscheweyh derivatives, in *Current Topics in Analytic Function Theory*, H.M. Srivastava and S. Owa (Eds.), pp. 13–28, World Scientific Publishing Company, Singapore, New Jersey, London and Hong Kong, 1992.
- [2] O.P. AHUJA AND M. NUNOKAWA, Neighborhoods of analytic functions defined by Ruscheweyh derivatives, *Math. Japon.*, **51** (2003), 487–492.
- [3] O. ALTINTAŞ AND S. OWA, Neighborhoods of certain analytic functions with negative coefficients, *Internat. J. Math. and Math. Sci.*, **19** (1996), 797–800.
- [4] O. ALTINTAŞ, Ö. ÖZKAN AND H.M. SRIVASTAVA, Neighborhoods of a class of analytic functions with negative coefficients, *Appl. Math. Lett.*, **13**(3) (2000), 63–67.
- [5] O. ALTINTAŞ, Ö. ÖZKAN AND H.M. SRIVASTAVA, Majorization by starlike functions of complex order, *Complex Variables Theory Appl.*, **46** (2001), 207–218.
- [6] O. ALTINTAŞ, Ö. ÖZKAN AND H.M. SRIVASTAVA, Neighborhoods of a certain family of multivalent functions with negative coefficients, *Comput. Math. Appl.*, **47** (2004) (to appear).
- [7] O. ALTINTAŞ AND H.M. SRIVASTAVA, Some majorization problems associated with p -valently starlike and convex functions of complex order, *East Asian Math. J.*, **17** (2001), 175–183.



Neighborhoods of Certain Classes of Analytic Functions of Complex Order

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents




Go Back

Close

Quit

Page 16 of 18

- [8] P.L. DUREN, *Univalent Functions*, A Series of Comprehensive Studies in Mathematics, Vol. **259**, Springer-Verlag, New York, Berlin, Heidelberg, and Tokyo, 1983.
- [9] A.W. GOODMAN, Univalent functions and nonanalytic curves, *Proc. Amer. Math. Soc.*, **8** (1957), 598–601.
- [10] J.-L. LIU AND H.M. SRIVASTAVA, Classes of meromorphically multivalent functions associated with the generalized hypergeometric function, *Math. Comput. Modelling*, **39** (2004), 21–34.
- [11] J.-L. LIU AND H.M. SRIVASTAVA, Subclasses of meromorphically multivalent functions associated with a certain linear operator, *Math. Comput. Modelling*, **39** (2004), 35–44.
- [12] M.A. NASR AND M.K. AOUF, Starlike function of complex order, *J. Natur. Sci. Math.*, **25** (1985), 1–12.
- [13] S. RUSCHEWEYH, New criteria for univalent functions, *Proc. Amer. Math. Soc.*, **49** (1975), 109–115.
- [14] S. RUSCHEWEYH, Neighborhoods of univalent functions, *Proc. Amer. Math. Soc.*, **81** (1981), 521–527.
- [15] H. SILVERMAN, Univalent functions with negative coefficients, *Proc. Amer. Math. Soc.*, **51** (1975), 109–116.
- [16] H. SILVERMAN, Neighborhoods of classes of analytic functions, *Far East J. Math. Sci.*, **3** (1995), 165–169.



**Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order**

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents




Go Back

Close

Quit

Page **17** of **18**

- [17] H.M. SRIVASTAVA AND S. OWA (Eds.), *Current Topics in Analytic Function Theory*, World Scientific Publishing Company, Singapore, New Jersey, London and Hong Kong, 1992.
- [18] P. WIATROWSKI, On the coefficients of some family of holomorphic functions, *Zeszyty Nauk. Uniw. Łódź Nauk. Mat.-Przyrod.* (2), **39** (1970), 75–85.



**Neighborhoods of Certain
Classes of Analytic Functions
of Complex Order**

G. Murugusundaramoorthy and
H.M. Srivastava

Title Page

Contents



Go Back

Close

Quit

Page 18 of 18